

APP955Hu01 100µg

Active Homeobox Protein C11 (HOXC11)

Organism Species: *Homo sapiens* (Human)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Met1~Leu304

Tags: N-terminal His-tag

Purity: >90%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Buffer Formulation: PBS, pH7.4, containing 0.01% Sarcosyl, 5%Trehalose .

Original Concentration: 200µg/mL

Applications: Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.7

Predicted Molecular Mass: 37.5kDa

Accurate Molecular Mass: 40kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

MFNSVNLGNFCSPSRKERGADFGERGSCASNLYLPSTYYMPEFSTVSSFLPQAPSRQISYPYSA
QVPPVREVSYGLEPSGKWHHRNSYSSCYAAADELMHRECLPPSTVTEILMKNEGSYGGHHHPS
APHATPAGFYSSVNKNSVLPQAFDRFFDNAYCGGGDPPAEPSCSGKGEAKGEPEAPPASGLASR
AEAGAEAEAEENTNPSSSGSAHSVAKEPAKGAAPNAPRTRKKRCPYSKFKIRELEREFFFNVIYN
KEKRLQLSRMLNLTDRQVKIWFQNRMMKEKKLSRDLQYFSGNPLL

[ACTIVITY]

Homeobox protein Hox-C11 (HOXC11) is a key regulatory protein. Encoded by the HOXC11 gene, it's a member of the homeobox transcription factor family. It functions importantly in embryonic morphogenesis and organogenesis, governing the spatial and temporal development of tissues and organs by binding to specific DNA sequences and regulating gene expression. Besides, Huntingtin (HTT) has been identified as an interactor of HOXC11, thus a functional binding ELISA assay was conducted to detect the interaction of recombinant human HOXC11 and recombinant human HTT. Briefly, biotin-linked HOXC11 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to HTT-coated microtiter wells and incubated for 1h at 37 °C. Wells were washed with PBST 3 times and incubation with Streptavidin-HRP for 30min, then wells were aspirated and washed 5 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37 °C. Finally, add 50 μ l stop solution to the wells and read at 450nm immediately. The binding activity of HOXC11 and HTT was shown in Figure 1, the EC₅₀ for this effect is 0.085ug/mL.



**Figure 1. The binding activity of recombinant human HOXC11 and recombinant human
HTT**

[IDENTIFICATION]




Figure 2. Gene Sequencing (extract)




Figure 3. SDS-PAGE

Sample: Active recombinant HOXC11, Human

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.