

**APA130Hu01 100µg**

**Active Tissue Inhibitors Of Metalloproteinase 4 (TIMP4)**

**Organism Species: *Homo sapiens (Human)***

***Instruction manual***

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

---

1st Edition (Apr, 2016)

## **[ PROPERTIES ]**

**Source:** Prokaryotic expression.

**Host:** *E. coli*

**Residues:** Cys32~Pro224

**Tags:** N-terminal His-tag

**Purity:** >92%

**Endotoxin Level:** <1.0EU per 1µg (determined by the LAL method).

**Buffer Formulation:** 20mM Tris, 150mM NaCl, pH8.0, containing 0.05% sarcosyl and 5% trehalose.

**Applications:** Cell culture; Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 7.2

**Predicted Molecular Mass:** 23.5kDa

**Accurate Molecular Mass:** 24kDa as determined by SDS-PAGE reducing conditions.

## **[ USAGE ]**

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## **[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


## **[ SEQUENCE ]**

```
                CAPAHPQQH ICHSALVIRA
KISSEKVVPA SADPADTEKM LRYEIKQIKM FKGFEKVKDV QYIYTPFDSS
LCGVKLEANS QKQYLLTGQV LSDGKVFHIL CNYIEPWEDL SLVQRESLNH
HYHLNCGCQI TTCYTVPCTI SAPNECLWTD WLLERKLYGY QAQHYVCMKH
VDGTCSWYRG HLPLRKEFVD IVQP
```

## **[ ACTIVITY ]**

Tissue Inhibitors Of Metalloproteinase 4 (TIMP4) is an enzyme that in humans is encoded by the TIMP4 gene. This gene belongs to the tissue inhibitor of metalloproteinases gene family. The proteins encoded by this gene family are inhibitors of the matrix metalloproteinases, a group of peptidases involved in degradation of the extracellular matrix. The secreted, netrin domain-containing protein encoded by this gene is involved in regulation of platelet aggregation and recruitment and may play role in hormonal regulation and endometrial tissue remodeling. Besides, Matrix Metalloproteinase 2 (MMP2) has been identified as an interactor of TIMP4, thus a binding ELISA assay was conducted to detect the interaction of recombinant human TIMP4 and recombinant human MMP2. Briefly, TIMP4 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100µL were then transferred to MMP2-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-TIMP4 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µL


stop solution to the wells and read at 450nm immediately. The binding activity of TIMP4 and MMP2 was shown in Figure 1, and this effect was in a dose dependent manner.




**Figure 1. The binding activity of TIMP4 with MMP2.**

**[ IDENTIFICATION ]**

TTGGCCGCGGCGCACCTCAGCAGCAGCCTGCGACCTGGCAGCTTGTGATTGGCCGAAATCTCCAGTGGAGAGGTGTTCCGGCCAGTGCAGACCTGCTGACACTGAAAAATCTCCGATGAAATCAACAGATAAAGATGTCCAAAGGTTTGGAGAAAGTCAAGGHTGTTCATATATCTATAAGCCCTTGGACTCTCCCTC  
CAPAHFQQQHICRSALVIRAKISSEKVVFPASADPADTEKMLRYEIKQIKMFKGFEEKYKDVQYIYTFDSSL




**Figure 2. Gene Sequencing (extract)**



**Figure 3. SDS-PAGE**

**Sample: Active recombinant TIMP4, Human**



**Figure 4. Western Blot**

**Sample: Recombinant TIMP4, Human;**

**Antibody: Rabbit Anti-Human TIMP4 Ab (PAA130Hu01)**

### **[ IMPORTANT NOTE ]**

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.