



CAB932Mi22

Anti-Glyceraldehyde-3-Phosphate Dehydrogenase (GAPDH)

Monoclonal Antibody

Organism Species: Multiple species

Instruction manual

FOR RESEARCH USE ONLY

3rd Edition (Revised in Sep, 2019)

[PROPERTIES]

Host: Mouse

Antibody isotype: IgG1 Kappa

Purification: Protein A/G Affinity Chromatography

Clone number: D2F4

Traits: Liquid

Concentration: 1mg/mL

Species reactivity: Human, Mouse, Rat, Guinea pig, Rabbit, Porcine, Bovine, Caprine, Ovine, Gallus, Equine

UOM: 100µl

Applications: Loading Control of WB.

[IMMUNOGEN]

Immunogen: Recombinant GAPDH (Gly2~Ser148) expressed in *E.coli*.

Accession No.: RPB932Hu01

[APPLICATIONS]

Western blotting: 1/10000-1/10000000 (0.1ng/ml-100ng/ml)

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.


Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the antibody at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under

appropriate storage condition.

[IDENTIFICATION]



Western Blot analysis of GAPDH in various cell lines

- Lane1: DU145 whole cell lysate
- Lane2: OS-RC-2 whole cell lysate
- Lane3: T-47D whole cell lysate
- Lane4: HEC-1B whole cell lysate
- Lane5: HepG2 whole cell lysate
- Lane6: HEK-293 whole cell lysate

Lysates/proteins at 20µg per lane.

Primary Ab: anti-GAPDH antibody (CAB932Mi22) at 0.1ng/ml

Secondary Ab: HRP-conjugated Rabbit anti-mouse antibody (SAA544Mu09) at 1/10000 dilution


Developed using the ECL technique.

Calculated MW: 36KD

Observed MW: 36KD

Exposure time: 1 min

Western Blot analysis of GAPDH in various tissues of different species



Lane1: Rat heart, whole tissue lysate

Lane2: Porcine brain, whole tissue lysate

Lane3: Bovine kidney, whole tissue lysate

Lane4: Guinea pig liver, whole tissue lysate

Lane5: Rabbit pancreas, whole tissue lysate

Lane6: Caprine spleen, whole tissue lysate

Lysates/proteins at 20 μ g per lane.

Primary Ab: anti-GAPDH antibody (CAB932Mi22) at 0.3ng/ml

Secondary Ab: HRP-conjugated Rabbit anti-mouse antibody (SAA544Mu09) at 1/10000 dilution

Developed using the ECL technique

Calculated MW: 36KD

Observed MW: 36KD

Exposure time: 1 min