
#### PAB180Rb51 Polyclonal Antibody to Clusterin (CLU) **Organism Species: Oryctolagus cuniculus (Rabbit)** Instruction manual

#### FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES



9th Edition (Revised in Jul, 2013)

-27kDa

kDa 75

50

37

25

20

15

10

Sample: Recombinant CLU, Rabbit

#### [IMMUNOGEN INFORMATION]

Immunogen: Recombinant CLU (Asn226~Glu447) expressed in E.coli.

Accession No.: RPB180Rb01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHHSGSEF-NIMPL SLYGPLNFQD MFQPFFEMIH QAQQAMDVHL HSPAYQTPNV EFITGGPDDR AVCKEIRHNS TGCLRMKDQC AKCQEILSVD CSANNPSQNQ LRQELNDSLR LAEELTKRYN ELLQSYQWKM LNTSSLLDQP NEQFNWVSQL ANLTQGPDQY YLRVSTVTSH TSESEAPSRV TEVVVKLFDS DPITITIPEE VSRDNPKFME TVAEKALQEY RKKKRVE

## [ANTIBODY SPECIFITY]

The antibody is a cavia polyclonal antibody raised against CLU. It has been selected for its ability to recognize CLU in immunohistochemical staining and western blotting.

### [APPLICATIONS]

Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500 Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

## [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant CLU (Asn226~Glu447) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$  0.02%.

# [<u>STORAGE</u>]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.