

#### PAP169Rb51

Polyclonal Antibody to Retinol Binding Protein 5, Cellular (RBP5)

Organism Species: Oryctolagus cuniculus (Rabbit)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

## [ PRODUCT INFORMATION ]

Immunogen: RBP5, Rabbit

Clonality: Polyclonal

Host: Cavia


Immunoglobulin Type: IgG

**Purification:** Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

**UOM**: 100µg



Sample: Recombinant RBP5, Rabbit

## [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant RBP5 (Glu19~Leu201) expressed in E.coli.

Accession No.: RPP169Rb01

**Sequence:** The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGHHHHHHSG SEF-ER DCRVSSFRVK ENFDKARFAG TWYAMAKKDP EGLFLQDNIV
AEFSVDENGH MSATAKGRVR LLNNWDVCAD MVGTFTDTED PAKFKMKYWG
VASFLQRGND DHWIIDTDYD TFAVQYSCRL LNFDGTCADS YSFVFSRDPH GLPPDVQKLV
ROROFFI CLS ROYRLIVHNG YCDDKSVRNLI



## [ANTIBODY SPECIFITY]

The antibody is a cavia polyclonal antibody raised against RBP5. It has been selected for its ability to recognize RBP5 in immunohistochemical staining and western blotting.

# [APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

# [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant RBP5 (Glu19~Leu201) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

## [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.