

LAB689Hu76 Biotin-Linked Monoclonal Antibody to Interleukin 28A (IL28A) Organism Species: Homo sapiens (Human) *Instruction manual* 

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

1304 Langham Creek Dr. Suite 226, Houston, TX 77064, USA | 001-888-960-7402 | www.cloud-chure.us | mailig/cloud-clone.as Export Processing Zone, Wuhan, Hubei 430056, PRC | 0086-000-880-0607 | www.cloud-clone.com | mailig/cloud-clone.com

# Cloud-Clone Corp.

# [PROPERTIES]

Source: Monoclonal antibody preparation Host: Mouse Purification: Protein A/G Affinity Chromatography. Clone number: C14 Label: Biotin Original Antibody: MAB689Hu26 Traits: Liquid Concentration: 500µg/mL UOM: 200µg Applications: WB; ICC; IHC-P; IHC-F; IF; ELISA.

# [IMMUNOGEN]

Immunogen: Recombinant IL28A (Val26~Val200) expressed in *E.coli*. Accession No.: RPB689Hu01

#### [APPLICATIONS]

Western blotting: 0.5-2ug/ml Immunocytochemistry in formalin fixed cells: 5-20ug/ml Immunohistochemistry in formalin fixed frozen section: 5-20ug/ml Immunohistochemistry in paraffin section: 5-20ug/ml Enzyme-linked Immunosorbent Assay: 0.05-2ug/ml Optimal working dilutions must be determined by end user.

# [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant IL28A disposed in loading buffer.

# Coud-Clone Corp.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH6.8), 1% SDS, 150mM NaCl, 50% glycerol, 0.02% BPB, 50mM DTT and 0.02% NaN<sub>3</sub>.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.