

LAA567Ra72
Biotin-Linked Monoclonal Antibody to S100 Calcium Binding Protein B (S100B)
Organism Species: Rattus norvegicus (Rat)

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

Instruction manual



# [PROPERTIES]

Source: Monoclonal antibody preparation

Host: Mouse

**Purification:** Protein A/G Affinity Chromatography.

Clone number: C6

Label: Biotin

Original Antibody: MAA567Ra21

Traits: Liquid

Concentration: 500µg/mL

**UOM**: 200µg

Applications: WB; IHC; ICC.

## [ IMMUNOGEN ]

Immunogen: Recombinant S100B (Met1~Glu92) expressed in E.coli.

Accession No.: RPA567Ra01

## [APPLICATIONS]

Western blotting: 0.5-2µg/mL

Immunohistochemistry: 5-20µg/mL Immunocytochemistry: 5-20µg/mL

Optimal working dilutions must be determined by end user.

## [FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing

0.05% Proclin-300, 50% glycerol.

## [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant S100B disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).



**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH6.8), 1% SDS, 150mM NaCl, 50% glycerol, 0.02% BPB, 50mM DTT and 0.02% NaN<sub>3</sub>.

#### [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.