EPB171Hu62 10μg

Eukaryotic Transferrin Receptor (TFR)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Eukaryotic expression

Host: 293F cell

Residues: Cys89~Phe760

Tags: N-terminal His Tag

Subcellular Location: Membrane, Secreted

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 5% Trehalose .

Original Concentration: 200μg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.8

Predicted Molecular Mass: 76.8kDa

Accurate Molecular Mass: 80kDa as determined by SDS-PAGE reducing conditions.

Phenomenon explanation:

The possible reasons that the actual band size differs from the predicted are as follows:

1.Splice variants: Alternative splicing may create different sized proteins from the same gene.

2. Relative charge: The composition of amino acids may affects the charge of the protein.

3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.

4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.

5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8ºC for one month.

Aliquot and store at -80ºC for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```
                                          CK GVEPKTECER
LAGTESPVRE EPGEDFPAAR RLYWDDLKRK LSEKLDSTDF TGTIKLLNEN
SYVPREAGSQ KDENLALYVE NQFREFKLSK VWRDQHFVKI QVKDSAQNSV
IIVDKNGRLV YLVENPGGYV AYSKAATVTG KLVHANFGTK KDFEDLYTPV
NGSIVIVRAG KITFAEKVAN AESLNAIGVL IYMDQTKFPI VNAELSFFGH
AHLGTGDPYT PGFPSFNHTQ FPPSRSSGLP NIPVQTISRA AAEKLFGNME
GDCPSDWKTD STCRMVTSES KNVKLTVSNV LKEIKILNIF GVIKGFVEPD
HYVVVGAQRD AWGPGAAKSG VGTALLLKLA QMFSDMVLKD GFQPSRSIIF
ASWSAGDFGS VGATEWLEGY LSSLHLKAFT YINLDKAVLG TSNFKVSASP
LLYTLIEKTM QNVKHPVTGQ FLYQDSNWAS KVEKLTLDNA AFPFLAYSGI
PAVSFCFCED TDYPYLGTTM DTYKELIERI PELNKVARAA AEVAGQFVIK
LTHDVELNLD YERYNSQLLS FVRDLNQYRA DIKEMGLSLQ WLYSARGDFF
RATSRLTTDF GNAEKTDRFV MKKLNDRVMR VEYHFLSPYV SPKESPFRHV
FWGSGSHTLP ALLENLKLRK QNNGAFNETL FRNQLALATW TIQGAANALS
GDVWDIDNEF
```

[IDENTIFICATION]



Figure . Gene Sequencing (extract)



Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.