

EPB367Hu61 10µg

Eukaryotic Vascular Endothelial Growth Factor Receptor 2 (VEGFR2)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Eukaryotic expression

Host: 293F cell

Residues: Ala20~Glu764

Tags: N-terminal His Tag

Subcellular Location: Membrane, Nucleus, Secreted, Cytoplasm, Endoplasmic reticulum lumen, Endosome

Purity: > 95%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 5% Trehalose.

Original Concentration: 50µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.1

Predicted Molecular Mass: 84.9kDa

Accurate Molecular Mass: 90kDa as determined by SDS-PAGE reducing conditions.

Phenomenon explanation:

The possible reasons that the actual band size differs from the predicted are as follows:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affect the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in ddH₂O to a concentration of 0.1-0.2 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[**SEQUENCE**]

```

                A SVGLPSVSLD LPRLSIQKDI LTIKANTTLQ
ITCRGQRDLG WLWPNNQSGS EQRVEVTECS DGLFCKTLTI PKVIGNDTGA
YKCFYRETDL ASVIYVYVQD YRSPFIASVS DQHGCVVYITE NKNKTVVIPC
LGSISNLNVS LCARYPEKRF VPDGNRISWD SKKGFTIPSY MISYAGMVFC
EAKINDESYQ SIMYIVVVVG YRIYDVVLSF SHGIELSVGE KLVLNCTART
ELNVGIDFNW EYPSSKHQHK KLVNRDLKTQ SGSEMKKFLS TLTIDGVTRS
DQGLYTCAAS SGLMTKKNST FVRVHEKPFV AFGSGMESLV EATVGERVRI
PAKYLGYPPE EIKWYKNGIP LESNHTIKAG HVLTIMEVSE RDTGNYTVIL
TNPISKEKQS HVVSLVVYVP PQIGEKSLIS PVDSYQYGTG QTLTCTVYAI
PPPHIHVYVW QLEEECANEP SQAVSVTNPY PCEEWRVSDV FQGGNKIEVN
KNQFALIEGK NKTVSTLVIQ AANVSALYKC EAVNKVGRGE RVISFHVTRG
PEITLQPDMQ PTEQESVSLW CTADRSTFEN LTWYKLGPPQ LPIHVGELPT
PVCKNLDTLW KLNATMFSNS TNDILIMELK NASLQDQGDY VCLAQDRKTK
KRHCVVRQLT VLERVAPTIT GNLENQTTSI GESIEVSCTA SGNPPPQIMW
FKDNETLVED SGIVLKDGNR NLTIRRVRKE DEGLYTCQAC SVLGCAKVEA
FFIIEGAQEK TNLE
```

[**IDENTIFICATION**]




Figure . Gene Sequencing (extract)




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.