

RPP389Hu01 100µg

Recombinant Fc Fragment Of IgG Binding Protein (FcgBP)

Organism Species: Homo sapiens (Human)

Instruction manual

66.2

45

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES.

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Cys5176~Ala5344

Tags: N-terminal His-Tag

Accession: Q9Y6R7

Host: E. coli

Subcellular Location: Secreted.

Purity: >95%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Formulation: Supplied as lyophilized form in 20mM Tris,

0.01% sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point: 8.3

Predicted Molecular Mass: 19.4kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in ddH₂O.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

CTHNG RYLPVNSSLL TSDCSERCSC SSSSGLTCQA AGCPPGRVCE VKAEARNCWA TRGLCVLSVG ANLTTFDGAR GATTSPGVYE LSSRCPGLQN TIPWYRVVAE VQICHGKTEA VGQVHIFFQD GMVTLTPNKG VWVNGLRVDL PAEKLASVSV SRTPDGSLLV RQKA

[REFERENCES]

- 1. Harada N., et al. (1997) J. Biol. Chem. 272:15232-15241.
- 2. Kobayashi K., et al. (2001) Immunol. Lett. 79:229-235.
- 3. O'Donovan N., et al. (2002) J. Endocrinol. 174:517-524.
- 4. Ramachandran P., et al. (2006) J. Proteome Res. 5:1493-1503.