RPA050Po01 10µg Recombinant Insulin Like Growth Factor 1 (IGF1) Organism Species: Sus scrofa; Porcine (Pig) *Instruction manual* 

### FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Coud-Clone Corp.

10th Edition (Revised in Jan, 2014)

#### [PROPERTIES] kDa 94 Residues: Gly49~Ala118 66.2 Tags: N-terminal GST-Tag 45 Accession: P16545 Host: E. coli 33 Subcellular Location: Secreted. 26 **Purity:** >90% Endotoxin Level: <1.0EU per 1µg 20 (determined by the LAL method). Formulation: Supplied as lyophilized form in 10mM 14.4 PBS, pH7.4, containing 1mM DTT, 5% trehalose, 0.01% sarcosyl and preservative. 15% SDS-PAGE Predicted isoelectric point: 5.9 Predicted Molecular Mass: 34.2kDa Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.)

## [ <u>USAGE</u> ]

Reconstitute in sterile ddH<sub>2</sub>O.

# Cloud-Clone Corp.

## [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

## [<u>SEQUENCES</u>]

The sequence of the target protein is listed below.

GP ETLCGAELVD ALQFVCGDRG FYFNKPTGYG SSSRRAPQTG IVDECCFRSC DLRRLEMYCA PLKPAKSA

## [REFERENCES]

- 1. Mueller M., Brem G. (1990) Nucleic Acids Res. 18:364-364.
- 2. Tavakkol A., et al. (1988) Mol. Endocrinol. 2:674-681.
- 3. Weller P.A., et al. (1993) J. Mol. Endocrinol. 11:201-211.
- 4. Gregoraszczuk E.L., et al. (2007) J. Reprod. Dev. 53:289-295.