

RPF552Hu02 100µg

Recombinant Lysyl Oxidase Like Protein 2 (LOXL2)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Ser544~Ser751

Tags: N-terminal His-Tag

Accession: Q9Y4K0

Host: *E. coli*

Subcellular Location: Secreted, extracellular space, extracellular matrix, basement membrane.

Nucleus. Chromosome.

Purity: >95%

Endotoxin Level: <1.0EU per 1µg
(determined by the LAL method).


Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 1mM DTT, 5% trehalose, 0.01% sarcosyl and preservative.

Predicted isoelectric point: 5.5

Predicted Molecular Mass: 25.6kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)



[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

SETAPDL VLNAEMVQQT TYLEDPRMFM LQCAMEENCL SASAAQTDPT TGYRLLRFS
SQIHNNQSD FRPKNGRHAW IWHDCRHYH SMEVFTHYDL LNLNGTKVAE
GHKASFLED TECEGDIQKN YECANFGDQG ITMGCWDMYR HDIDCQWVDI TDVPPGDYLF
QVVINPNFEV AESDYSNNIM KCRSRYDGHR IWMYNCHIGG S

[REFERENCES]

1. Saito H., *et al.* (1997) J. Biol. Chem. 272:8157-8160.
2. Jourdan-Le Saux C., *et al.* (1999) J. Biol. Chem. 274:12939-12944.
3. Peinado H., *et al.* (2005) EMBO J. 24:3446-3458.
4. Schietke R., *et al.* (2010) J. Biol. Chem. 285:6658-6669.