

MAA782Si22

Monoclonal Antibody to Carbonic Anhydrase II (CA2) Organism Species: Rhesus monkey (Simian) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES.

9th Edition (Revised in Jul, 2013)

←31kDa

Western Blot

kDa 75

50

37

25 20

15

10

[PRODUCT INFORMATION]

Immunogen: CA2, Simian

Clonality: Monoclonal Clone number: C6

Host: Mouse

Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 500µg/mL

Sample: Recombinant CA2, Simian

UOM: 200µg [IMMUNOGEN INFORMATION]

Immunogen: Recombinant CA2 (Met1~Lys260) expressed in E.coli.

Accession No.: RPA782Si01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF- MSHHWGYGKH NGPEHWHKDF PIAKGQRQSP VDIDTHTAKY DPSLKPLSVS YDQATSLRIL NNGHSFNVEF DDSQDKAVIK GGPLDGTYRL IQFHFHWGSL DGQGSEHTVD KKKYAAELHL VHWNTKYGDF GKAVQQPDGL AVLGIFLKVG SAKPGLOKVV DVLDSIKTKG KSADFTNFDP RGLLPESLDY WTYPGSLTTP PLLECVTWIV I KEPISVSSE OMSKERKI NE NGEGEPEELM VDNWRPAOPL KNROIKASEK



[ANTIBODY SPECIFITY]

The antibody is a mouse monoclonal antibody raised against CA2. It has been selected for its ability to recognize CA2 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.