

MAF331Hu21 Monoclonal Antibody to Cellular Retinoic Acid Binding Protein 2 (CRABP2) Organism Species: *Homo sapiens (Human)* Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

Cond-Clone Corp.

[PROPERTIES]

Source: Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG2b Kappa

Purification: Protein A + Protein G affinity chromatography

Clone number: C1

Traits: Liquid

Concentration: 1mg/ml

UOM: 100µl

Cross Reactivity: Rat; Porcine

Applications: WB

[IMMUNOGEN]

Immunogen: Recombinant CRABP2 (Met1~Glu138) expressed in E.coli

Accession No.: RPF331Hu01

[APPLICATIONS]

Western blotting: 0.01-2µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300,

50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the

Contend Cloud - Clone Corp.

expiration date under appropriate storage condition.

[IDENTIFICATION]





Western Blot; Sample: Lane1: Rat Skin Western Blot; Sample: Lane1: Porcine lysate; Lane2: Porcine Skin lysate; Esophagus lysate; Lane2: MCF7 cell Lane3: Porcine Esophagus lysate lysate Primary Ab: 0.2µg/ml Mouse Anti-Primary Ab: 0.2µg/ml Mouse Anti-Human CRABP2 Antibody Second Ab: Human CRABP2 Antibody 0.2µg/mL HRP-Linked Caprine Anti-Second Ab: 0.2µg/mL HRP-Linked Mouse IgG Polyclonal Antibody Caprine Anti-Mouse IgG Polyclonal (Catalog: SAA544Mu19) Antibody (Catalog: SAA544Mu19)

[<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.