

MAB995Hu24

Monoclonal Antibody to Hemojuvelin (HJV)
Organism Species: Homo sapiens (Human) *Instruction manual* 

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



### [PROPERTIES]

**Source:** Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG2b Kappa

**Purification:** Protein A/G Affinity Chromatography.

Clone number: C6

Traits: Liquid

Concentration: 500µg/mL

**UOM**: 200µg

Applications: WB; IHC; ICC; IP.

# [ IMMUNOGEN ]

Immunogen: Recombinant HJV (Lys234~Ser416) expressed in E.coli.

Accession No.: RPB995Hu01

### [APPLICATIONS]

Western blotting: 0.5-5µg/mL

Immunohistochemistry: 5-30µg/mL Immunocytochemistry: 5-30µg/mL

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing

0.05% Proclin-300, 50% glycerol.

# [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant HJV disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine (DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control.

Not used for other purposes.



**Loading Buffer:** 100mM Tris(pH6.8), 1% SDS, 150mM NaCl, 50% glycerol, 0.02% BPB, 50mM DTT and 0.02% NaN<sub>3</sub>.

## [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]




Figure 1. Western Blot

A. Sample: Recombinant HJV, Human

B. Lane1: Porcine Liver Tissue

Lane2: Rat Liver Tissue

Primary Ab: 2µg/mL Mouse Anti-Human HJV Ab
Second Ab: 1:5000 Dilution of HRP-Linked Rabbit
Anti-Mouse IgG Ab (Catalog: SAA544Mu09)




Figure 2. DAB staining on IHC-P Samples: