

MAB934Hu26

Monoclonal Antibody to Leucine Rich Alpha-2-Glycoprotein 1 (LRG1)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

# Cond-Clone Corp.

## [PROPERTIES]

Source: Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG2a Kappa

Purification: Protein A + Protein G affinity chromatography

Clone number: D2

Traits: Liquid

Concentration: 1mg/ml

**UOM:** 100µl

Cross Reactivity: N/A

Applications: WB

[IMMUNOGEN]

Immunogen: Recombinant LRG1 (Thr37~Leu340) expressed in E.coli

Accession No.: RPB934Hu01

#### [APPLICATIONS]

Western blotting: 0.5-3µg/mL;

Immunohistochemistry: 5-30µg/mL;

Immunocytochemistry: 5-30µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined

## Cond-Clone Corp.

by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [IDENTIFICATION]






Figure. Western Blot; Sample: Recombinant LRG1, Human.





Western Blot; Sample: Human UrineWestern Blot; Sample: Human SerumPrimary Ab: 3µg/ml Mouse Anti-HumanPrimary Ab: 3µg/ml Mouse Anti-HumanLRG1 AntibodyLRG1 AntibodySecond Ab: 0.2µg/mL HRP-LinkedSecond Ab: 0.2µg/mL HRP-LinkedCaprine Anti-Mouse IgG PolyclonalCaprine Anti-Mouse IgG PolyclonalAntibodyAntibody(Catalog: SAA544Mu19)(Catalog: SAA544Mu19)

### [<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.