

**MAA105Hu23**

**Monoclonal Antibody to Nerve Growth Factor (NGF)**

**Organism Species: *Homo sapiens* (Human)**

***Instruction manual***

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

13th Edition (Revised in Aug, 2023)

**[ PROPERTIES ]**

**Source:** Monoclonal antibody preparation

**Host:** Mouse

**Antibody isotype:** IgG1 Kappa

**Purification:** Protein A + Protein G affinity chromatography

**Clone number:** C2

**Traits:** Liquid

**Concentration:** 1mg/ml

**UOM:** 100µl

**Cross Reactivity:** Porcine; Bovine.

**Applications:** WB

**[ IMMUNOGEN ]**

**Immunogen:** Recombinant NGF (Glu19~Arg239) expressed in *E.coli*

**Accession No.:** RPA105Hu01

**[ APPLICATIONS ]**

Western blotting: 0.01-2µg/mL;

Optimal working dilutions must be determined by end user.

**[ FORMULATION ]**

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

**[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the

expiration date under appropriate storage condition.

**[ IDENTIFICATION ]**





Figure. Western Blot; Sample: Recombinant NGF, Human.



Western Blot; Sample: Lane1: Porcine Spleen lysate; Lane2: Porcine Cerebrum lysate; Lane3: Porcine Skin lysate; Lane4: Porcine Pancreas lysate; Lane5: Bovine Spleen lysate; Lane6: Bovine Cerebrum lysate; Lane7: U87MG cell lysate  
Primary Ab: 0.5?g/ml Mouse Anti-Human NGF Antibody  
Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Mouse IgG Polyclonal Antibody  
(Catalog: SAA544Mu19)

**[ IMPORTANT NOTE ]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.