

**MAA837Hu22** 

Monoclonal Antibody to Nitric Oxide Synthase 2, Inducible (NOS2)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



# [PROPERTIES]

Source: Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG2a Lambda

**Purification:** Protein A + Protein G affinity chromatography

Clone number: C1

Traits: Liquid

Concentration: 1mg/mL

**UOM:** 100µL

Cross Reactivity: Mouse

Applications: WB; IHC

### [ IMMUNOGEN ]

Immunogen: Recombinant NOS2 (Thr734~Gly966) expressed in E.coli

Accession No.: RPA837Hu02

#### [ APPLICATIONS ]

Western blotting: 0.01-3µg/mL;

Immunohistochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no


# Cloud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


## [ IDENTIFICATION ]



Western Blot; Sample: Human Serum
Primary Ab: 3µg/ml Mouse Anti-Human
NOS2 Antibody Second Ab: 0.2µg/ml
HRP-Linked Caprine Anti-Mouse IgG
Polyclonal Antibody (Catalog:
SAA544Mu19)




DAB staining on IHC-P; Samples:
Human Liver Tissue; Primary Ab:
10μg/ml Mouse Anti-Human NOS2
Antibody Second Ab: 2μg/mL HRPLinked Caprine Anti-Mouse IgG
Polyclonal Antibody (Catalog:
SAA544Mu19)



DAB staining on IHC-P;

Sample: Human Liver Tissue
Primary Ab: 10µg/ml Mouse AntiHuman NOS2 Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2µg/ml HRP-Linked
Caprine Anti-Mouse IgG Polyclonal
Antibody
(Catalog: SAA544Mu19)



Western blot; Samples: lysates of RAW264.7 cell line treated with 1  $\mu$ M staurosporine (STS) for the indicated times.

Primary Ab: 0.3ug/ml Mouse Anti-Human NOS2 Antibody Second Ab: 0.2µg/mL HRP-Linked



Caprine Anti-Mouse IgG Polyclonal
Antibody
(Catalog: SAA544Mu19)

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.