



P95741Hu01 Phosphoglucomutase 5 (PGM5) **Organism: Homo sapiens (Human)** Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

15% SDS-PAGE

Human PGM5

[DESCRIPTION]

 SDS-PAGE reducing conditions. Formulation: Supplied as lyophilized form in PBS, pH 7.4, containing 5% sucrose, 0.01% sarcosyl. Endotoxin Level: <1.0 EU per 1µg (determined by the LAL method). Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.) 	Protein Names: Phosphoglucomutase 5	Human PGI
Species: Human66.2Size: 100µg45Source: Escherichia coli-derived33Subcellular Location: Cell junction, adherens26junction. Cytoplasm, cytoskeleton.20 [PROPERTIES] 20Residues: Leu182~Gly296 (Accession # Q15124), with N-terminal His-Tag.14.4Grade & Purity: >95%, 16kDa as determined by SDS-PAGE reducing conditions.15% SDS-FFormulation: Supplied as lyophilized form in PBS, pH15% SDS-F7.4, containing 5% sucrose, 0.01% sarcosyl.Endotoxin Level: <1.0 EU per 1µg (determined by the LAL method).Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.)	Synonyms: PGM5, PGMRP	
Source: Escherichia coli-derived 33 Subcellular Location: Cell junction, adherens 33 junction. Cytoplasm, cytoskeleton. 20 [PROPERTIES] 20 Residues: Leu182~Gly296 (Accession # Q15124), 14.4 with N-terminal His-Tag. 14.4 Grade & Purity: >95%, 16kDa as determined by 15% SDS-F Formulation: Supplied as lyophilized form in PBS, pH 15% SDS-F Formulation: Supplied as lyophilized form in PBS, pH 1.4 7.4, containing 5% sucrose, 0.01% sarcosyl. Endotoxin Level: <1.0 EU per 1µg (determined by	Species: Human	
Subcellular Location: Cell junction, adherens33junction. Cytoplasm, cytoskeleton.20[PROPERTIES]20Residues: Leu182~Gly296 (Accession # Q15124), with N-terminal His-Tag.14.4Grade & Purity: >95%, 16kDa as determined by SDS-PAGE reducing conditions.15% SDS-FFormulation: Supplied as lyophilized form in PBS, pH15% SDS-F7.4, containing 5% sucrose, 0.01% sarcosyl.Endotoxin Level: <1.0 EU per 1µg (determined by the LAL method).Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.)	Size: 100µg	45
Subcellular Location: Cell junction, adherens junction. Cytoplasm, cytoskeleton.26[PROPERTIES]20Residues: Leu182~Gly296 (Accession # Q15124), with N-terminal His-Tag.14.4Grade & Purity: >95%, 16kDa as determined by SDS-PAGE reducing conditions.15% SDS-FFormulation: Supplied as lyophilized form in PBS, pH15% SDS-F7.4, containing 5% sucrose, 0.01% sarcosyl.Endotoxin Level: <1.0 EU per 1µg (determined by the LAL method).Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.)	Source: Escherichia coli-derived	33
junction. Cytoplasm, cytoskeleton. [PROPERTIES] 20 Residues: Leu182~Gly296 (Accession # Q15124), 14.4 with N-terminal His-Tag. Grade & Purity: >95%, 16kDa as determined by SDS-PAGE reducing conditions. Formulation: Supplied as lyophilized form in PBS, pH 7.4, containing 5% sucrose, 0.01% sarcosyl. Endotoxin Level: <1.0 EU per 1µg (determined by the LAL method). Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.)	Subcellular Location: Cell junction, adherens	
Residues: Leu182~Gly296 (Accession # Q15124), 14.4 with N-terminal His-Tag. 14.4 Grade & Purity: >95%, 16kDa as determined by 15% SDS-PAGE reducing conditions. SDS-PAGE reducing conditions. 15% SDS-F Formulation: Supplied as lyophilized form in PBS, pH 15% SDS-F 7.4, containing 5% sucrose, 0.01% sarcosyl. Endotoxin Level: <1.0 EU per 1µg (determined by the LAL method).	junction. Cytoplasm, cytoskeleton.	20
with N-terminal His-Tag. 14.4 Grade & Purity: >95%, 16kDa as determined by 15% SDS-PAGE reducing conditions. SDS-PAGE reducing conditions. 15% SDS-F Formulation: Supplied as lyophilized form in PBS, pH 15% SDS-F 7.4, containing 5% sucrose, 0.01% sarcosyl. Endotoxin Level: <1.0 EU per 1µg (determined by the LAL method).	[PROPERTIES]	20
 with N-terminal His-Tag. Grade & Purity: >95%, 16kDa as determined by SDS-PAGE reducing conditions. Formulation: Supplied as lyophilized form in PBS, pH 7.4, containing 5% sucrose, 0.01% sarcosyl. Endotoxin Level: <1.0 EU per 1µg (determined by the LAL method). Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.) 	Residues: Leu182~Gly296 (Accession # Q15124),	14.4
SDS-PAGE reducing conditions.15% SDS-FFormulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% sucrose, 0.01% sarcosyl.Endotoxin Level: <1.0 EU per 1µg (determined by the LAL method).Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.)	with N-terminal His-Tag.	14.4
 SDS-PAGE reducing conditions. Formulation: Supplied as lyophilized form in PBS, pH 7.4, containing 5% sucrose, 0.01% sarcosyl. Endotoxin Level: <1.0 EU per 1µg (determined by the LAL method). Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.) 	Grade & Purity: >95%, 16kDa as determined by	
 7.4, containing 5% sucrose, 0.01% sarcosyl. Endotoxin Level: <1.0 EU per 1µg (determined by the LAL method). Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.) 	SDS-PAGE reducing conditions.	15% SDS-P/
 Endotoxin Level: <1.0 EU per 1µg (determined by the LAL method). Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.) 	Formulation: Supplied as lyophilized form in PBS, pH	
the LAL method). Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.)	7.4, containing 5% sucrose, 0.01% sarcosyl.	
Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.)	Endotoxin Level: <1.0 EU per 1µg (determined by	
(May be suitable for use in other assays to be determined by the end user.)	the LAL method).	
	Applications: SDS-PAGE; WB; ELISA; IP.	
Predicted Malacular Massa 44 40	(May be suitable for use in other assays to be determined by	y the end user.)
Predicted Molecular Mass: 14.1kDa	Predicted Molecular Mass: 14.1kDa	

Predicted isoelectric point: 5.8

Unique product Superb quality Client favorite Nicest service @ ISO9001:2008; @ ISO13485:2003; C \in





[PREPARATION]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below. MGHHHHHHSGSEF- LENKFKPFR VEIVDPVDIY LNLLRTIFDF HAIKGLLTGP SQLKIRIDAM HGVMGPYVRK VLCDELGAPA NSAINCVPLE DFGGQHPDPN LTYATTLLEA MKGGEYGFGA AFDADG

Unique product Superb quality Client favorite Nicest service @ ISO9001:2008; @ ISO13485:2003; \in