

PAE892Hu01**Polyclonal Antibody to A Kinase Anchor Protein 12 (AKAP12)****Organism Species: Homo sapiens (Human)****Instruction manual**

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]**Immunogen:** AKAP12, Human**Clonality:** Polyclonal**Host:** Rabbit**Immunoglobulin Type:** IgG**Purification:** Affinity Chromatography.**Applications:** WB, ICC, IHC-P, IHC-F, ELISA**Concentration:** 200µg/mL**UOM:** 100µg**[IMMUNOGEN INFORMATION]****Immunogen:** Recombinant AKAP12 (Pro1485~Ser1782) expressed in *E.coli*.**Accession No.:** RPE892Hu01**Sequence:** The target protein is fused with two N-terminal Tags, His-tag and T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- PVIVSA TTKKGLSSDL
EGEKTTSKWK KSDEVDEQVA CQEVKVSVAI EDLEPENGIL ELETKSSKLV QNIIQTAVDQ
FVRTEETATE MLTSELQTQA HVIKADSQDA GQETEKEGEE PQASAQDETP ITSAKEESES
TAVGQAHSKI SKDMSEASEK TMTVEVEGST VNDQQLEEVV LPSEEEGGGA GTKSVPEDDG
HALLAERIEK SLVEPKEDDK GDDVDDPENQ NSALADTDAS GGLTKESPDT NGPKQKEKED
AQEVELQEGK VHSSEDKAIT PQAQEELQKQ ERESAKSELT ES

[ANTIBODY SPECIFICITY]

The antibody is a rabbit polyclonal antibody raised against AKAP12. It has been selected for its ability to recognize AKAP12 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500


Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100


Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[**IMAGES**]



Used in DAB staining on formalin fixed
paraffin-embedded ovary tissue



Used in Western Blot, Sample:
Recombinant AKAP12, Human

[**CONTENTS**]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[**QUALITY CONTROL**]

Content: The quality control contains recombinant AKAP12 (Pro1485~Ser1782) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.
5uL per well when used in enhanced chemiluminescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[**STORAGE**]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.