

# PAA816Po01 Polyclonal Antibody to Alpha-1-Acid Glycoprotein (a1AGP) Organism Species: Sus scrofa; Porcine (Pig) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

## [ PRODUCT INFORMATION ]

Immunogen: a1AGP, Pig

Clonality: Polyclonal

Host: Rabbit

Immunoglobulin Type: IgG


Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

**UOM:** 100µg

#### 9th Edition (Revised in Jul, 2013)



Sample: Recombinant a1AGP, Pig

#### [ IMMUNOGEN INFORMATION ]

**Immunogen:** Recombinant a1AGP (Leu2~Glu171) linked with GKGR expressed in *E.coli*.

Accession No.: RPA816Po01

**Sequence:** The target protein is fused with two N-terminal Tags, His-tag and T7-tag, its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- LCANLTAVP ITNATLDLIS GKWYYIGSAF RNPQYNESAR SIQAAFFFFD PKPAEDKINL REYQTIGNQC IYNDSSLKVH RENGSLSKHE MGREHVADLL LTKVPKTFML INSLHDKNNV GLSFYADKAE VTPEQMKEFH DAIECTGIHK SEITYTDEKK DLCGPLEKQH E GKGR



#### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against a1AGP. It has been selected for its ability to recognize a1AGP in immunohistochemical staining and western blotting.

#### [APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

#### [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

#### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant a1AGP disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

### [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.