

PAB254Mu01

Polyclonal Antibody to Angiostatin (ANG)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

Cond-Clone Corp.

[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.73mg/ml

UOM: 100µl

Cross Reactivity: Human; Rat

Applications: WB; IHC; ICC; IP.

[IMMUNOGEN]

Immunogen: Recombinant ANG (Val98~Trp436) expressed in E.coli

Accession No.: RPB254Mu01

[APPLICATIONS]

Western blotting: 0.5-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN3, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

Cloud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]		
Western Blot KDa 70 44 33 26 22 18 14 10	kDa 100- 70- 44- 33- 27- 22-	kDa 100— 70— 44—
Figure. Western Blot; Sample:	Western Blot; Sample: Mouse Heart	Western Blot; Sample: Mouse Lung
Recombinant ANG, Mouse.	lysate;	lysate;
	Primary Ab: 2µg/ml Rabbit Anti-Mouse	Primary Ab: 2µg/ml Rabbit Anti-Mouse
	ANG Antibody	ANG Antibody
	Second Ab: 0.2µg/mL HRP-Linked	Second Ab: 0.2µg/mL HRP-Linked
	Caprine Anti-Rabbit IgG Polyclonal	Caprine Anti-Rabbit IgG Polyclonal
	Antibody	Antibody
	(Catalog: SAA544Rb19)	(Catalog: SAA544Rb19)

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.