

# PAA519Rb51 Polyclonal Antibody to Apolipoprotein A1 (APOA1) Organism Species: Oryctolagus cuniculus (Rabbit) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

#### [ PRODUCT INFORMATION ]

Immunogen: APOA1, Rabbit

Clonality: Polyclonal

Host: Cavia


Immunoglobulin Type: IgG

**Purification:** Affinity Chromatography.


Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

**UOM**: 100µg



Western Blot



Sample: Recombinant APOA1, Rabbit

#### [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant APOA1 (Asp25~Gln265) expressed in E.coli.

Accession No.: RPA519Rb01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGS-DEPRSS WDKIKDFATV YVDTVKDSGR EYVAQFEASA FGKQLNLKLL DNWDSLSSTV SKLQEQLGPV TQEFWDNLEK ETEGLREEMN KDLQEVRQKV QPFLDEFQKK WQEEVERYRQ KVEPLGAELG ESARQKLTEL QEKLSPLAEE LRDSARTHVD TLRTKLAPYS NELQRLAARL ESIKEGGGAK LAEYQAKARE HLSVLSEKAR PALEDLRQGL LPVLESFKAS VQNVVDEATK KLNTQ



#### [ANTIBODY SPECIFITY]

The antibody is a cavia polyclonal antibody raised against APOA1. It has been selected for its ability to recognize APOA1 in immunohistochemical staining and western blotting.

## [APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

## [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant APOA1 (Asp25~Gln265) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

## [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.