PAA896Po01 Polyclonal Antibody to Cystatin 3 (CST3) Organism Species: Sus scrofa; Porcine (Pig) *Instruction manual* 

# FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

1304 Langham Creek Dr. Soite 226, Houston, TX 77094, USA | 001-888-960-7402 | www.choud-choue.ox | mail@choud-chone.as Export Processing Zone, Wuhan, Hubei 436056, PRC | 0086-000-880-0687 | www.choud-clone.com | mail@choud-chone.com

# Cloud-Clone Corp.

#### [PROPERTIES]

Source: Polyclonal antibody preparation Host: Rabbit Purification: Antigen-specific Affinity Chromatography. Traits: Liquid Concentration: 200µg/mL UOM: 100µg Applications: WB; ICC; IHC-P; IHC-F; ELISA; IP; IF; FCM.

#### [IMMUNOGEN]

Immunogen: Recombinant CST3 (Ala42~Val127) expressed in *E.coli*. Accession No.: RPA896Po01

#### [APPLICATIONS]

Western blotting: 0.5-2ug/ml Immunocytochemistry in formalin fixed cells: 5-20ug/ml Immunohistochemistry in formalin fixed frozen section: 5-20ug/ml Immunohistochemistry in paraffin section: 5-20ug/ml Enzyme-linked Immunosorbent Assay: 0.05-2ug/ml Optimal working dilutions must be determined by end user.

## [FORMULATION]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were

# Coud-Clone Corp.

observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [IDENTIFICATION]




Figure 1. Western Blot A. Sample: Recombinant CST3, Porcine