

#### PAC903Mu01 Polyclonal Antibody to Deiodinase, Iodothyronine, Type II (DIO2) **Organism Species: Mus musculus (Mouse)** Instruction manual

#### FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES



9th Edition (Revised in Jul, 2013)



### [IMMUNOGEN INFORMATION]

Immunogen: Recombinant DIO2 (Ser30~Phe259) expressed in E.coli.

Accession No.: RPC903Mu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and MBP-tag and its sequence is listed below.

MKIEEGKLVI WINGDKGYNG LAEVGKKFEK DTGIKVTVEH PDKLEEKFPQ VAATGDGPDI IFWAHDRFGG YAQSGLLAEI TPDKAFQDKL YPFTWDAVRY NGKLIAYPIA VEALSLIYNK DLLPNPPKTW EEIPALDKEL KAKGKSALMF NLQEPYFTWP LIAADGGYAF KYENGKYDIK DVGVDNAGAK AGLTFLVDLI KNKHMNADTD YSIAEAAFNK GETAMTINGP WAWSNIDTSK VNYGVTVLPT FKGQPSKPFV GVLSAGINAA SPNKELAKEF LENYLLTDEG LEAVNKDKPL GAVALKSYEE ELAKDPRIAA TMENAQKGEI MPNIPOMSAF WYAVRTAVIN AASGRQTVDE ALKDAQTGST SGSGHHHHHH SAGLVPRGST AIGMKETAAA KFEROHMDSP DLGTLEVLFO GPLGSEF- S VILLKHVALL LSRSKSTRGE WRRMLTSEGL RCVWNSFLLD AYKOVKLGED APNSSVVHVS NPESGNNYAS EKTADGAECH LLDFASAERP LVVNFGSATU PPFTRQLPAF

RQLVEEFSSV ADFLLVYIDE AHPSDGWAVP GDSSLSFEVK KHRNQEDRCA AAHQLLERFS LPPQCQVVAD RMDNNANVAY GVAFERVCIV QRRKIAYLGG KGPFSYNLQE VRSWLEKNF

## [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against DIO2. It has been selected for its ability to recognize DIO2 in immunohistochemical staining and western blotting.

### [APPLICATIONS]

Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500 Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

## [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## [ QUALITY CONTROL ]

**Content:** The quality control is from mouse brain extract known to contain the target protein DIO2, disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$  0.02%.

# [ <u>STORAGE</u> ]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.