

PAA512Mu01

Polyclonal Antibody to Fibrinogen Like Protein 2 (FGL2)
Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: FGL2, Mouse **Purification:** Affinity Chromatography.

Clonality: Polyclonal Applications: WB, ICC, IHC-P, IHC-F, ELISA

Host: Rabbit Concentration: 200µg/mL

Immunoglobulin Type: IgG **UOM**: 100μg

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant FGL2 (Val20~Pro432) expressed in E.coli.

Accession No.: RPA512Mu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQM GRGSEF- V EEHNLTEGLE DASAQAACPA RLEGSGRCEG SQCPFQLTLP TLTIQLPRQL GSMEEVLKEV RTLKEAVDSL KKSCQDCKLQ ADDHRDPGGN GGNGAETAED SRVQELESQV NKLSSELKNA KDQIQGLQGR LETLHLVNMN NIENYVDNKV ANLTVVVNSL DGKCSKCPSQ EHMQSQPVQH LIYKDCSDHY VLGRRSSGAY RVTPDHRNSS FEVYCDMETM GGGWTVLQAR LDGSTNFTRE WKDYKAGFGN LEREFWLGND KIHLLTKSKE MILRIDLEDF NGLTLYALYD QFYVANEFLK YRLHIGNYNG TAGDALRFSR HYNHDLRFFT TPDRDNDRYP SGNCGLYYSS GWWFDSCLSA NLNGKYYHQK YKGVRNGIFW GTWPGINQAQ PGGYKSSFKQ AKMMIRPKNF KP

[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against FGL2. It has been selected for its ability to recognize FGL2 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400


Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500


Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[IMAGES]



Used in DAB staining on fromalin fixed paraffin-embedded Spleen tissue



Used in Western Blot, Sample:
Recombinant FGL2, Mouse

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant FGL2 (Val20~Pro432) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.