

PAB494Hu01

Polyclonal Antibody to Fibromodulin (FMOD)
Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: FMOD, Human **Purification:** Affinity Chromatography.

Clonality: Polyclonal Applications: WB, ICC, IHC-P, IHC-F, ELISA

Host: Rabbit Concentration: 200µg/mL

Immunoglobulin Type: IgG **UOM**: 100μg

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant FMOD (Gln19~Ile376) expressed in E.coli.

Accession No.: RPB494Hu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- QY EDDPHWWFHY LRSQQSTYYD PYDPYPYETY EPYPYGVDEG PAYTYGSPSP PDPRDCPQEC DCPPNFPTAM YCDNRNLKYL PFVPSRMKYV YFQNNQITSI QEGVFDNATG LLWIALHGNQ ITSDKVGRKV FSKLRHLERL YLDHNNLTRM PGPLPRSLRE LHLDHNQISR VPNNALEGLE NLTALYLQHN EIQEVGSSMR GLRSLILLDL SYNHLRKVPD GLPSALEQLY MEHNNVYTVP DSYFRGAPKL LYVRLSHNSL TNNGLASNTF NSSSLLELDL SYNQLQKIPP VNTNLENLYL QGNRINEFSI SSFCTVVDVV NFSKLQVLRL DGNEIKRSAM PADAPLCLRL ASLIEI

[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against FMOD. It has been selected for its ability to recognize FMOD in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400




Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500


Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[IMAGES]



Used in DAB staining on fromalin fixed paraffin-embedded Liver tissue



Used in Western Blot, Sample:

Recombinant FMOD, Human

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant FMOD (Gln19~Ile376) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.