

PAJ048Ra01 Polyclonal Antibody to Gastrokine 2 (GKN2) Organism Species: *Rattus norvegicus (Rat) Instruction manual*

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

Cond-Clone Corp.

[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 1mg/ml

UOM: 100µg(100µl)

Applications: WB; IHC; ICC; IP.

[<u>IMMUNOGEN</u>]

Immunogen: Recombinant GKN2 (Lys21~Leu184 (Accession # Q29TV8)) expressed in E.coli

Accession No.: RPJ048Ra01

[APPLICATIONS]

Western blotting: 0.5-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN3, 50%

glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the

Cloud-Clone Corp.

expiration date under appropriate storage condition.

[IDENTIFICATION]








Figure. Western Blot; Sample: Recombinant GKN2, Rat.

Western Blot; Sample: Mouse StomachWestern Blot; Sample: Rat StomachIysate;Iysate;Primary Ab: 1µg/ml Rabbit Anti-RatPrimary Ab: 1µg/ml Rabbit Anti-RatGKN2 AntibodyGKN2 AntibodySecond Ab: 0.2µg/mL HRP-LinkedSecond Ab: 0.2µg/mL HRP-LinkedCaprine Anti-Rabbit IgG PolyclonalCaprine Anti-Rabbit IgG PolyclonalAntibodyAntibody(Catalog: SAA544Rb19)(Catalog: SAA544Rb19)

[<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.