

PAA068Mu03 Polyclonal Antibody to Glial Fibrillary Acidic Protein (GFAP) Organism Species: *Mus musculus (Mouse) Instruction manual*

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

Cond-Clone Corp.

[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 1mg/ml

UOM: 100µg(100µl)

Cross Reactivity: Human; Rat

Applications: WB; IHC; ICC; IP.

[<u>IMMUNOGEN</u>]

Immunogen: Recombinant GFAP (Lys108~Gln211 (Accession # P03995)) expressed in E.coli

Accession No.: RPA068Mu03

[APPLICATIONS]

Western blotting: 0.5-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 12 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

Cloud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]








Figure. Western Blot; Sample: Recombinant GFAP, Mouse.

Western Blot; Sample: MouseDAB staining on IHC-P;Cerebrum lysate;Samples: Mouse Brain Tissue;Primary Ab: 2µg/mL Rabbit Anti-Mouse Primary Ab: 10µg/ml Rabbit Anti-MouseGFAP AntibodyGFAP AntibodySecond Ab: 0.2µg/mL HRP-LinkedSecond Ab: 2µg/mL HRP-LinkedCaprine Anti-Rabbit IgG PolyclonalCaprine Anti-Rabbit IgG PolyclonalAntibodyAntibody(Catalog: SAA544Rb19)(Catalog: SAA544Rb19)

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.