

P	Δ	C ₁	6	6	R	h	5	1

Polyclonal Antibody to HLA Class II Histocompatibility Antigen, DR Alpha Chain (HLA-DRA)

Organism Species: Oryctolagus cuniculus (Rabbit)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Guinea pig

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/ml

UOM: 200µl

Cross Reactivity: N/A

Applications: WB; IHC.

[IMMUNOGEN]

Immunogen: Recombinant HLA-DRA (Ala26~Cys221+LEHHHHHHH) expressed in E.coli

Accession No.: RPC166Rb01

[APPLICATIONS]

Western blotting: 0.01-2µg/mL

Immunohistochemistry: 5-20µg/mL

Immunocytochemistry: 5-20µg/mL

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

Coud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]




Figure. Western Blot; Sample: Recombinant MHCDRa, Rabbit.





Figure. Western Blot; Sample: Lane1:
Rabbit Spleen Tissue; Lane2: Rabbit
Lymph Tissue; Lane3: Rabbit Thymus
Tissue.



DAB staining on IHC-P; Samples:
Rabbit Spleen Tissue; Primary Ab:
10µg/ml Cavia Anti-Rabbit MHCDRa
Antibody Second Ab: 2µg/mL HRPLinked Rabbit Anti-Cavia IgG
Polyclonal Antibody (Catalog:
SAA544Gu09)

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.