

PAB953Mu01

Polyclonal Antibody

To Inhibitor Of Nuclear Factor Kappa B Kinase Interacting Protein (IKBIP)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

### [ PRODUCT INFORMATION ]

Immunogen: IKBIP, Mouse Purification: Affinity Chromatography.

Clonality: Polyclonal Applications: WB, ICC, IHC-P, IHC-F, ELISA

Host: Rabbit Concentration: 200µg/mL

**Immunoglobulin Type**: IgG **UOM**: 100μg

#### [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant IKBIP (Thr78~Arg301) expressed in E.coli.

Accession No.: RPB953Mu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- TES SEFQGLQSKI SLISSKYQEC EALVEQLKAF QIAAHLKLLQ EEIHEMKTWS NRITERQGTL NNTLTRLSED IIKVDQGTAS MAKDMGLKIT SVKTDVRRIS GLVTEVESLT DAVQALGNKV KKVETATVEN IGDLLSSSID RTSALRKTAS ENAKRIDSVA QRLAELQGDF DEHTDRFLSL ESDRAKVLKA VSFANDLKPK VSNLKKDFSR LEPLVDDLTL R

#### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against IKBIP. It has been selected for its ability to recognize IKBIP in immunohistochemical staining and western blotting.

### [APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500




Immunohistochemistry in formalin fixed frozen section: 1:100-500


Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

# [ IMAGES ]



Used in DAB staining on fromalin fixed paraffin-embedded Kidney tissue



Used in Western Blot, Sample:
Recombinant IKBIP, Mouse

# [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant IKBIP (Thr78~Arg301) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

### [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.