Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

[PRODUCT INFORMATION] Western Blot kDa 75 Immunogen: LAT2, Human 50 **Clonality:** Polyclonal 37 Host: Rabbit 25 Immunoglobulin Type: IgG 20 **Purification:** Affinity Chromatography. 15 Applications: WB, ICC, IHC-P, IHC-F, ELISA 10 Concentration: 200µg/mL **UOM:** 100µg Sample: Recombinant LAT2, Human

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant LAT2 (Arg30~Met209) expressed in E.coli. Accession No.: RPP555Hu01 Sequence: The target protein is fused with two N-terminal Tags, His-tag and T7-tag and its sequence is listed below. MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- R PGAKRSEKIY QQRSLREDQQ SFTGSRTYSL VGQAWPGPLA DMAPTRKDKL LQFYPSLEDP

ASSRYQNFSK GSRHGSEEAY IDPIAMEYYN WGRFSKPPED DDANSYENVL ICKOKTTETG AQQEGIGGLC RGDLSLSLAL KTGPTSGLCP SASPEEDEES EDYQNSASIH QWRESRKVM

9th Edition (Revised in Jul, 2013)

←24kDa

[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against LAT2. It has been selected for its ability to recognize LAT2 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500 Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant LAT2 (Arg30~Met209) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$ 0.02%.

[<u>STORAGE</u>]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.