

PAB954Mu01

Polyclonal Antibody to Monoamine Oxidase A (MAOA)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



### [PROPERTIES]

**Source:** Polyclonal antibody preparation

Host: Rabbit

**Purification:** Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/ml

**UOM**: 20µl

Cross Reactivity: Human; Rat

Applications: WB; IHC; ICC; IP.

#### [ IMMUNOGEN ]

Immunogen: Recombinant MAOA (Gln229~Gly428) expressed in E.coli

Accession No.: RPB954Mu01

### [ APPLICATIONS ]

Western blotting: 0.01-2µg/mL

Immunohistochemistry: 5-30µg/mL

Immunocytochemistry: 5-30µg/mL

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN3, 50% glycerol.

#### [STORAGE AND STABILITY]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

# Coud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]






Figure. Western Blot; Sample: Recombinant MAOA, Mouse.



DAB staining on IHC-P; Samples: Mouse Liver Tissue;


Primary Ab: 30µg/ml Rabbit Anti-Mouse Primary Ab: 30µg/ml Rabbit Anti-Mouse




DAB staining on IHC-P; Samples: Mouse Spleen Tissue;

MAOA Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)


MAOA Antibody
Second Ab: 2μg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



DAB staining on IHC-P;



DAB staining on IHC-P;



DAB staining on IHC-P; Samples: Mouse Uterus Tissue;

Samples: Mouse Stomach Tissue;

Samples: Mouse Cerebrum Tissue;

Primary Ab: 30μg/ml Rabbit Anti-Mouse Primary Ab: 30μg/ml Rabbit Anti-Mouse Primary Ab: 30μg/ml Rabbit Anti-Mouse


MAOA Antibody
Second Ab: 2μg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody

(Catalog: SAA544Rb19)

MAOA Antibody
Second Ab: 2μg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)

MAOA Antibody
Second Ab: 2μg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)

# Coud-Clone Corp.



FITC staining on IF;

Samples: Human HepG2 Cells;

Primary Ab: 10µg/ml Rabbit Anti-Mouse

MAOA Antibody

Second Ab: 1µg/ml FITC-Linked

Caprine Anti-Rabbit IgG Polyclonal

Antibody

(Catalog: SAA544Rb18)

## [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.