PAA484Ra02

Polyclonal Antibody to N-Terminal Pro-Atrial Natriuretic Peptide (NT-ProANP) Organism Species: Rattus norvegicus (Rat) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

1304 Langham Creek Dr. Soite 226, Houston, TX 77094, USA | 001-888-960-7402 | www.choud-choue.ox | mail@choud-chone.as Export Processing Zone, Wuhan, Hubei 436056, PRC | 0086-000-880-0687 | www.choud-clone.com | mail@choud-chone.com

Cloud-Clone Corp.

[PROPERTIES]

Source: Polyclonal antibody preparation Host: Rabbit Purification: Antigen-specific Affinity Chromatography. Traits: Liquid Concentration: 200µg/mL UOM: 100µg Applications: WB; IHC; ICC; IP.

[IMMUNOGEN]

Immunogen: Recombinant NT-ProANP (His27~His134) expressed in *E.coli*. **Accession No.:** RPA484Ra02

[APPLICATIONS]

Western blotting: 0.5-2µg/mL Immunocytochemistry in formalin fixed cells: 5-20µg/mL Immunohistochemistry in formalin fixed frozen section: 5-20µg/mL Immunohistochemistry in paraffin section: 5-20µg/mL Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant NT-ProANP disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH6.8), 1% SDS, 150mM NaCl, 50% glycerol, 0.02% BPB, 50mM DTT and 0.02% NaN₃.

Coud-Clone Corp.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]





Figure 1. Western Blot A. Sample: Recombinant NT-ProANP, Rat B. Lane1: Rat Heart Tissue Lane2: Mouse Heart Tissue Primary Ab: 1µg/mL Rabbit Anti-Rat NT-ProANP Ab Second Ab: 1:2000 Dilution of HRP-Linked Guinea pig Anti-Rabbit Ab (Catalog: SAA544Rb59)

Coud-Clone Corp.






Figure 2. DAB staining on IHC-P

Samples:

- A. Rat Skeletal Muscle Tissue
- B. Rat Stomach Tissue
- C. Rat Intestine Tissue

1304 Langham Creek Dr. Saire 226, Houston, TX 77084, USA | 001-888-960-7402 | www.cloud-cloue.ox | mail@cloud-close.ax Export Processing Zone, Wuhan, Hubei 430056, PRC | 0086-008-0687 | www.cloud-close.com | mail@cloud-close.com