

PAL915Hu01 Polyclonal Antibody to Nei Endonuclease VIII Like Protein 1 (NEIL1) Organism Species: Homo sapiens (Human) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

# Cloud-Clone Corp.

### [PROPERTIES]

Source: Polyclonal antibody preparation Host: Rabbit Purification: Antigen-specific Affinity Chromatography. Traits: Liquid Concentration: 200µg/mL UOM: 100µg Applications: WB; IHC; ICC; IP.

#### [IMMUNOGEN]

Immunogen: Recombinant NEIL1 (Ser43~Ala314) expressed in *E.coli*. Accession No.: RPL915Hu01

### [APPLICATIONS]

Western blotting: 0.5-2µg/mL Immunohistochemistry: 5-20µg/mL Immunocytochemistry: 5-20µg/mL Optimal working dilutions must be determined by end user.

### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

### [QUALITY CONTROL]

**Content:** The quality control contains recombinant NEIL1 disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control.

Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH6.8), 1% SDS, 150mM NaCl, 50% glycerol, 0.02% BPB, 50mM DTT and 0.02% NaN<sub>3</sub>.



#### [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

#### [IDENTIFICATION]






Figure 1. Western Blot Sample: Recombinant NEIL1, Human

# Cloud-Clone Corp.





# Figure 2. DAB staining on IHC-P

#### Samples:

- A. Human Lung Cancer Tissue
- B. Human Liver Tissue
- C. Human Stomach Tissue