

PAB083Mu01

Polyclonal Antibody to Superoxide Dismutase 2, Mitochondrial (SOD2)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



## [PROPERTIES]

**Source:** Polyclonal antibody preparation

Host: Rabbit

**Purification:** Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.34mg/ml

**UOM**: 100µl

Cross Reactivity: Human; Rat

Applications: WB; IHC; ICC; IP.

#### [ IMMUNOGEN ]

Immunogen: Recombinant SOD2 (Lys25~Lys222 (Accession # P09671)) expressed in E.coli

Accession No.: RPB083Mu01

### [ APPLICATIONS ]

Western blotting: 0.5-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

# Coud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]




Figure. Western Blot; Sample: Recombinant SOD2, Mouse.





Figure. Western Blot; Sample: Lane1: Mouse Heart Tissue; Lane2: Mouse



DAB staining on IHC-P; Samples: Mouse Stomach Tissue; Liver Tissue; Lane3: Rat Brain Tissue. Primary Ab: 10µg/ml Rabbit Anti-Mouse

SOD2 Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

(Catalog: SAA544Rb19)

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.