

PAA525Hu02

Polyclonal Antibody to Tissue Plasminogen Activator (tPA)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



## [PROPERTIES]

**Source:** Polyclonal antibody preparation

Host: Rabbit

**Purification:** Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/ml

**UOM**: 20µl

Cross Reactivity: Mouse; Rat

Applications: WB; IHC; ICC; IP.

#### [ IMMUNOGEN ]

Immunogen: Recombinant tPA (Ile311~Pro562) expressed in E.coli

Accession No.: RPA525Hu02

### [ APPLICATIONS ]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN3, 50% glycerol.

#### [STORAGE AND STABILITY]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

# Coud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

# [ IDENTIFICATION ]






Figure.Western Blot;Sample: Recombinant tPA, Human.



DAB staining on IHC-P; Samples: Human Kidney Tissue; Primary Ab: 10µg/ml Rabbit Anti-Human tPA Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody


(Catalog: SAA544Rb19)



DAB staining on IHC-P; Samples: Human Prostate cancer Tissue: Primary Ab: 10µg/ml Rabbit Anti-Human tPA Antibody Second Ab: 2µg/mL HRP-Linked


Caprine Anti-Rabbit IgG Polyclonal Antibody

(Catalog: SAA544Rb19)




DAB staining on IHC-P;

Samples: Human Breast cancer Tissue; Samples: Human Skin cancer Tissue; Primary Ab: 10µg/ml Rabbit Anti-Human tPA Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



DAB staining on IHC-P;

Primary Ab: 10µg/ml Rabbit Anti-Human tPA Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



DAB staining on IHC-P;

Samples: Human Liver cancer Tissue; Primary Ab: 10µg/ml Rabbit Anti-Human tPA Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)

# Coud-Clone Corp.




Figure. Western Blot; Sample. Lane1:

Human Mcf7 Cells; Lane2: Mouse

Pancreas Tissue; Lane3: Mouse Kidney

Tissue.

## [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.