

#### PAB989Mu01

Polyclonal Antibody to Toll Like Receptor 3 (TLR3) Organism Species: *Mus musculus (Mouse)* 

Instruction manual

#### FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

# Cond-Clone Corp.

## [PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/ml

**UOM:** 50µl

Cross Reactivity: Rat

Applications: WB; IHC; ICC; IP.

#### [<u>IMMUNOGEN</u>]

Immunogen: Recombinant TLR3 (Gly574~Met713 (Accession # Q99MB1)) expressed in E.coli

Accession No.: RPB989Mu01

### [APPLICATIONS]

Western blotting: 0.01-2µg/mL

Immunohistochemistry: 5-20µg/mL

Immunocytochemistry: 5-20µg/mL

Optimal working dilutions must be determined by end user.

### [FORMULATION]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN3, 50% glycerol.

### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.


**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

# Cloud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

# [IDENTIFICATION]





Western Blot; Sample: Lane1: Mouse Western Blot; Sample: Lane1: A549 cell Lung lysate; Lane2: Mouse Liver lysate lysate; Lane2: Rat Liver lysate Primary Ab: 1µg/ml Rabbit Anti-Mouse Primary Ab: 1µg/ml Rabbit Anti-Mouse TLR3 Antibody Second Ab: 0.2µg/mL TLR3 Antibody HRP-Linked Caprine Anti-Rabbit IgG Second Ab: 0.2µg/mL HRP-Linked Polyclonal Antibody (Catalog: SAA544Rb19) Antibody (Catalog: SAA544Rb19)

# [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.