

CSI049Ca01

Primary Canine Microglia Cells (MC)

Organism Species: Canis familiaris; Canine (Dog)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

1st Edition (Revised in Feb, 2024)

## [ DESCRIPTION ]

Cell Type: Microglia Cell

Synonyms: MC

Species: Canis familiaris; Canine (Dog)

Tissue Source: Cerebral cortex

Size: >5×10<sup>5</sup>cell/vial

## [PROPERTIES]

Cell activity: >85% (Viability by Trypan Blue Exclusion).

Formulation: Frozen 1 mL or T25 flask.

Biosafety: Negative for HIV-1, HBV, HCV, mycoplasma, bacteria, yeast and fungi.

Applications: For research use only. It is not approved for human or animal use, or for application in

clinical diagnostic procedures. **Growth Properties:** Adherent

## [CONTENTS]

Form & Buffer: Supplied as solution form in frozen stock solution, containing 90% FBS+10% DMSO.

## [USAGE]

Upon receiving the cells in a T-25 flask at room temperature, immediately transfer the cells to 37°C, 5% incubator; the cells in vials, directly and immediately transfer the cells from dry ice to liquid nitrogen.

#### **Culture conditions:**

DMEM/F12+5%FBS+1% Microglia Cell Growth Supplement+1%Penicillin-Streptomycin Solution

Temperature: 37°C

Condition: 95% air, 5% carbon dioxide

### Cell recovery:

After receiving the cells, shake at 37°C in a water bath until completely dissolved, transfer to a 15 ml centrifuge tube, add 3-5 times complete culture solution, 1000 rpm for 5 min, discard the supernatant, and place in a T25 flask for culture.

#### **Maintenance Culture:**

Change the medium to fresh supplemented medium the next morning after establishing a culture from cryopreserved cells. Change the medium every two to three days thereafter.

## [Shipping]

Dry ice.


## [STORAGE]

Upon receiving, directly and immediately transfer the cells from dry ice to liquid nitrogen and keep the cells in liquid nitrogen until they are needed for experiments.

## [ IMPORTANT NOTE ]

- 1. It is not recommended that Primary Canine Microglia Cells (MC) be subcultured beyond their initial plating.
- 2. The cell is for research use only, and we will not be responsible for any issue if the cell was used in clinical diagnostic or any other procedures.

# [Figure]






Figure 2

Figure 1

- Figure 1 Morphology of Primary Canine Microglia Cells (100X)
- Figure 2 Morphology of Primary Canine Microglia Cells (200X)