

RPC271Hu02 10ug

Recombinant 5-Hydroxytryptamine Receptor 1A (HTR1A)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Met1~Gln422

Tags: His and TrxA Taq

Subcellular Location: Membrane

Purity: > 80%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% skl, 5%Trehalose.

Original Concentration: 80µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.1

Predicted Molecular Mass: 66.5kDa

Accurate Molecular Mass: 70kDa as determined by SDS-PAGE reducing conditions.

Phenomenon explanation:

The possible reasons that the actual band size differs from the predicted are as follows:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in ddH₂O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.



Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```
MDVLSPGQGN NTTSPPAPFE TGGNTTGISD VTVSYQVITS LLLGTLIFCA VLGNACVVAA IALERSLQNV ANYLIGSLAV TDLMVSVLVL PMAALYQVLN KWTLGQVTCD LFIALDVLCC TSSILHLCAI ALDRYWAITD PIDYVNKRTP RRAAALISLT WLIGFLISIP PMLGWRTPED RSDPDACTIS KDHGYTIYST FGAFYIPLLL MLVLYGRIFR AARFRIRKTV KKVEKTGADT RHGASPAPQP KKSVNGESGS RNWRLGVESK AGGALCANGA VRQGDDGAAL EVIEVHRVGN SKEHLPLPSE AGPTPCAPAS FERKNERNAE AKRKMALARE RKTVKTLGII MGTFILCWLP FFIVALVLPF CESSCHMPTL LGAIINWLGY SNSLLNPVIY AYFNKDFQNA FKKIIKCKFC RQ
```

[IDENTIFICATION]




Figure . Gene Sequencing (extract)






Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.