

RPA620Mu01 10µg

Recombinant A Disintegrin And Metalloprotease 8 (ADAM8)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



### [ PROPERTIES ]

**Source:** Prokaryotic expression

Host: E.coli

Residues: Glu145~Cys493

Tags: N-terminal His Tag

**Subcellular Location:** Membrane

**Purity:** > 95%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.3

Predicted Molecular Mass: 42.4kDa

Accurate Molecular Mass: 44kDa as determined by SDS-PAGE reducing conditions.

#### [USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



## [SEQUENCE]

				EEGQHA
MYQAKHLQQK	AGTCGVKDTN	LNDLGPRALE	IYRAQPRNWL	IPRETRYVEL
YVVADSQEFQ	KLGSREAVRQ	RVLEVVNHVD	KLYQELSFRV	VLVGLEIWNK
DKFYISRYAN	VTLENFLSWR	EQNLQGQHPH	DNVQLITGVD	FIGSTVGLAK
VSALCSRHSG	AVNQDHSKNS	IGVASTMAHE	LGHNLGMSHD	EDIPGCYCPE
PREGGGCIMT	ESIGSKFPRI	FSRCSKIDLE	SFVTKPQTGC	LTNVPDVNRF
VGGPVCGNLF	VEHGEQCDCG	TPQDCQNPCC	NATTCQLVKG	AECASGTCCH
ECKVKPAGEV	CRLSKDKCDL	EEFCDGRKPT	<b>CPEDAFOONG</b>	TPC

## [ IDENTIFICATION ]




Figure. SDS-PAGE

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.