

RPG385Hu01 10µg

Recombinant ATPase, Cu⁺⁺ Transporting Alpha Polypeptide (ATP7a)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Ser9~Leu444

Tags: N-terminal His Tag

Subcellular Location: Membrane, Golgi apparatus

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose .

Original Concentration: 100µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.8

Predicted Molecular Mass: 51.0kDa

Accurate Molecular Mass: 51kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

SV TISVEGMTCN SCVWTIEQQI GKVNGVHHIK VSLEEKNATI IYDPKLQTP KTLQEAIIDM GFDAVIHNP
PLPVLDTLTF LVTASLTLP WDHIQSTLL KTKGVTDIKI YPQKRTVAVT IIPSIVNANQ IKELVPELSL DTGTLEKKS
GACEDHSMAQ AGEVVLKMKV EGMTCHSCTS TIEGKIGKLQ GVQRIKVS L DNQEATIVYQ PHLISVEEMK KQIEAMGFPA
FVKKQPKYLK LGAIDVERL KNTPVKSSEG SQQRSPSYTN DSTATFIIDG MHCKSCVSNI ESTLSALQY VSSIVVLEN
RSAIVKYNAS SVTPESLRKA IEAVSPGLYR VSITSEVES TSNPSSSSL QKIPLNVVSQ PLTQETVINI DGMTCNQCVQ
SIEGVISKK PGVKSIRVSL ANSNGTVEYD PLLTSPETLR GAIEDMGFDA TL

[IDENTIFICATION]




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.