

RPQ709Hu01 100ug

Recombinant Activating Signal Cointegrator 1 Complex Subunit 3 (ASCC3)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



# [ PROPERTIES ]

**Source:** Prokaryotic expression

Host: E.coli

Residues: Pro1812~Tyr2176

Tags: N-terminal His Tag

Subcellular Location: Nucleus, Cytoplasm

**Purity:** > 95%

Traits: Freeze-dried powder

**Buffer formulation:** 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% skl, 5%Trehalose.

Original Concentration: 80µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 7.1

Predicted Molecular Mass: 45.5kDa

**Accurate Molecular Mass:** 46kDa as determined by SDS-PAGE reducing conditions.

#### [USAGE]

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ SEQUENCE ]



```
PLTYGRIAS YYYLKHQTVK MFKDRLKPEC STEELLSILS
DAEEYTDLPV RHNEDHMNSE LAKCLPIESN PHSFDSPHTK AHLLQAHLS
RAMLPCPDYD TDTKTVLDQA LRVCQAMLDV AANQGWLVTV LNITNLIQMV
IQGRWLKDSS LLTLPNIENH HLHLFKKWKP IMKGPHARGR TSIESLPELI
HACGGKDHVF SSMVESELHA AKTKQAWNFL SHLPVINVGI SVKGSWDDLV
EGHNELSVST LTADKRDDNK WIKLHADQEY VLQVSLQRVH FGFHKGKPES
CAVTPRFPKS KDEGWFLILG EVDKRELIAL KRVGYIRNHH VASLSFYTPE
IPGRYIYTLY FMSDCYLGLD QQYDIY
```

## [ IDENTIFICATION ]




Figure. SDS-PAGE

### [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.