

RPB015Mu01 50µg

Recombinant Alpha-1-Antichymotrypsin (a1ACT)

Organism Species: *Mus musculus* (Mouse)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Asp23~Arg253

Tags: N-terminal His Tag

Subcellular Location: Secreted

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 150µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.8

Predicted Molecular Mass: 29.9kDa

Accurate Molecular Mass: 30kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

DGTLGMDA AVQEDHDNGT QLDSLTLASI
 NTDFAFSLYK ELVLKNPDKN IVFSPLSISA ALAVMSLGAK GNTLEEILEG
 LKFNLTTETSE ADIHQGFQHL LQRLNQPKDQ VQISTGSALF IEKRQQILTE
 FQEKAKTLYQ AEAFTADFQQ PRQAKKLIND YVRKQTQGM I KELVSDLDKR
 TLMVLVNYIY FKAKWKVPFD PLDTFKSEFY AGKRRPVIVP MMSMEDLTP
 YFR

[IDENTIFICATION]

GGATCCGATGGCACGTTGGGAATGGATGCTGCACTCCAAAGAACCATGACAAATGGGACACAAC TGACAGTCTCACATTGGCCCTCCATCAACACTGACTTTGCTTCAGCCTCTACAAGGAAGCTGGT TTTGAAAGATCCAGATACAAAATTGTCTCTCCACCATTAGCATCTCAGGGC





Figure . Gene Sequencing (extract)



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.