

RPB967Ra01 200µg

Recombinant Apolipoprotein A4 (APOA4)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Glu21~Ser391

Tags: N-terminal His and GST Tag

Subcellular Location: Secreted

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: 100mMNaHCO₃, 500mMNaCl, pH8.3, containing 1mM EDTA, 1mM DTT,

0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.1

Predicted Molecular Mass: 72.4kDa

Accurate Molecular Mass: 72kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 100mM NaHCO3, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[SEQUENCE]

		EVTSDQVANV	MWDYFTQLSN	NAKEAVEQLQ	
KTDVTQQLNT	LFQDKLGNIN	TYADDLQNKL	VPFAVQLSGH	LTKETERVRE	
EIQKELEDLR	ANMMPHANKV	SQMFGDNVQK	LQEHLRPYAT	DLQAQINAQT	
QDMKRQLTPY	IQRMQTTIQD	NVENLQSSMV	PFANELKEKF	NQNMEGLKGQ	
LTPRANELKA	TIDQNLEDLR	SRLAPLAEGV	QEKLNHQMEG	LAFQMKKNAE	
ELQTKVSTNI	DQLQKNLAPL	VEDVQSKLKG	NTEGLQKSLE	DLNKQLDQQV	
EVFRRAVEPL	GDKFNMALVQ	QMEKFRQQLG	SDSGDVESHL	SFLEKNLREK	
VSSFMSTLOK	KGSPDOPLAL	PLPEOVOEOV	OEOVOPKPLE	S	

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.