

RPB214Mu01 200µg

Recombinant Aspartate Aminotransferase (AST)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Met1~Gln413

Tags: N-terminal His Tag

Subcellular Location: Cytoplasm

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% skl, 5%Trehalose.

Original Concentration: 150µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.9

Predicted Molecular Mass: 47.5kDa

Accurate Molecular Mass: 43kDa as determined by SDS-PAGE reducing conditions.

Phenomenon explanation:

The possible reasons that the actual band size differs from the predicted are as follows:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]



Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```
MAPPSVFAQV PQAPPVLVFK LTADFRDDPD PRKVNLGVGA YRTDESQPWV LPVVRKVEQK IANDNSLNHE YLPILGLAEF RSCASRLVLG DNSLAIRENR VGGVQSLGGT GALRIGADFL GRWYNGTDNK NTPIYVSSPT WENHNAVFSA AGFKDIRPYC YWDAEKRGLD LQGFLNDLEN APEFSIFVLH ACAHNPTGTD PTPEQWKQIA AVMQRRFLFP FFDSAYQGFA SGDLEKDAWA IRYFVSEGFE LFCAQSFSKN FGLYNERVGN LTVVGKESDS VLRVLSQMEK IVRITWSNPP AQGARIVAAT LSDPELFKEW KGNVKTMADR ILTMRSELRA RLEALKTPGT WSHITEQIGM FSFTGLNPKQ VEYLVNEKHI YLLPSGRINM CGLTTKNLDY VATSIHEAVT KIQ
```

[IDENTIFICATION]





Figure. Gene Sequencing (Extract)





[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.