RPA655Ra01 100µg

Recombinant Calcium/Calmodulin Dependent

Protein Kinase II Gamma (CAMK2g)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)


[PROPERTIES]


Residues: Tyr14~Lys301 Tags: Two N-terminal Tags, His-tag and T7-tag Accession: P11730 Host: E. coli Subcellular Location: Sarcoplasmic reticulum membrane, Peripheral membrane protein, Cytoplasmic side Inferred. **Purity: >95%** Endotoxin Level: <1.0EU per 1µg (determined by the LAL method). Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% trehalose, 0.01% sarcosyl. Predicted isoelectric point: 8.1 Predicted Molecular Mass: 36.7kDa Applications: SDS-PAGE; WB; ELISA; IP.

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

(May be suitable for use in other assays to be determined by the end user.)





15% SDS-PAGE

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[<u>SEQUENCES</u>]

The sequence of the target protein is listed below.

YQLFEEL GKGAFSVVRR CVKKTSTQEY AAKIINTKKL SARDHQKLER EARICRLLKH PNIVRLHDSI SEEGFHYLVF DLVTGGELFE DIVAREYYSE ADASHCIHQI LESVNHIHQH DIVHRDLKPE NLLLASKCKG AAVKLADFGL AIEVQGEQQA WFGFAGTPGY LSPEVLRKDP YGKPVDIWAC GVILYILLVG YPPFWDEDQH KLYQQIKAGA YDFPSPEWDT VTPEAKNLIN QMLTINPAKR ITADQALKHP WVCQRSTVAS MMHRQETVEC LRKFNARRKL K