

RPB180Ov01 1 Recombinant Clusterin (CLU) Organism Species: *Ovis aries; Ovine (Sheep)* Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

Coud-Clone Corp.

[PROPERTIES]

Source: Prokaryotic expression
Host: *E.coli*

Residues: Ile20~Glu439

Tags: N-terminal His Tag

Subcellular Location: Nucleus, Secreted

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 100µg

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.7

Predicted Molecular Mass: 52.6kDa

Accurate Molecular Mass: 53kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

Control Cloud-Clone Corp.

I SDKELQEMST EGSKYVNKEI KNALKEVKQI KTQIEQTNEE RKLLLSSLEE AKKKKEDALN DTRDSENKLK ASQGVCNETM TALWEECKPC LKQTCMKFYA RVCRSGSGLV GHQLEEFLNQ SSPFYFWING DRIDSLMEND REQSHVMDVM EDSFTRASSI MDELFQDRFF PRRPQDTQYY SPFSSFPRGS LFFNPKSRFA RNVMPFPLLE PLNFHDVFQP FYDMIRQAQQ AMDAHLQRTP YHFPVTEFTE NNDRTVCKEI RHNSTGCLRM KDQCEKCQEI LEVDCSASSP TQTLLRQQLN TSLQLAEKFS RLYDQLLQSY QQKMLDTSTL LKQLNEQFTW VSQLANLTQS DDQYYLQVFT VNSHSSDPSI PSGLTKVVVK LFNSFPITVT VPQEVSSPNF MENVAEKALQ QYRRKSHEE

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.