

RPB878Ca01 100µg Recombinant Coagulation Factor VIII (F8) Organism Species: *Canis familiaris; Canine (Dog) Instruction manual* 

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

# Coud-Clone Corp.

## [PROPERTIES]

Source: Prokaryotic expression Host: E.coli Residues: Met2191~Cys2337 Tags: N-terminal His Tag Subcellular Location: Secreted **Purity:** > 90% Traits: Freeze-dried powder Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300. Original Concentration: 200µg/mL Applications: Positive Control; Immunogen; SDS-PAGE; WB. (May be suitable for use in other assays to be determined by the end user.) Predicted isoelectric point: 10.1 Predicted Molecular Mass: 17.9kDa Accurate Molecular Mass: 21/18/17kDa as determined by SDS-PAGE reducing conditions. Phenomenon explanation: The possible reasons that the actual band size differs from the predicted are as follows: 1.Splice variants: Alternative splicing may create different sized proteins from the same gene. 2. Relative charge: The composition of amino acids may affects the charge of the protein. 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.

4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.

5. Polymerization of the target protein: Dimerization, multimerization etc.

### [ <u>USAGE</u> ]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

# Cond-Clone Corp.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.


Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

#### [ <u>SEQUENCE</u> ]

MESKAISDAQ ITASSYLSSM LATWSPSQAR LHLQGRTNAW RPQANNPKEW LQVDFRKTMK VTGITTQGVK SLLISMYVKE FLISSSQDGH NWTLFLQNGK VKVFQGNRDS STPVRNRLEP PLVARYVRLH PQSWAHHIAL RLEVLGC

### [IDENTIFICATION]



## [<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.