

RPA888Hu01 50µg

**Recombinant Complement Component 4 (C4)** 

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



### [ PROPERTIES ]

**Source:** Prokaryotic expression

Host: E.coli

Residues: Asn975~Arg1336

Tags: N-terminal His Tag

Subcellular Location: Secreted

**Purity:** > 95%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% Sarcosyl, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.8

Predicted Molecular Mass: 42.5kDa

Accurate Molecular Mass: 42kDa as determined by SDS-PAGE reducing conditions.

#### [USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.


**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



## [SEQUENCE]

		NSYVRV	TASDPLDTLG	SEGALSPGGV
ASLLRLPRGC	GEQTMIYLAP	TLAASRYLDK	TEQWSTLPPE	TKDHAVDLIQ
KGYMRIQQFR	KADGSYAAWL	SRDSSTWLTA	FVLKVLSLAQ	EQVGGSPEKL
QETSNWLLSQ	QQADGSFQDP	CPVLDRSMQG	GLVGNDETVA	LTAFVTIALH
HGLAVFQDEG	AEPLKQRVEA	SISKANSFLG	EKASAGLLGA	HAAAITAYAL
TLTKAPVDLL	GVAHNNLMAM	AQETGDNLYW	GSVTGSQSNA	VSPTPAPRNP
SDPMPQAPAL	WIETTAYALL	HLLLHEGKAE	MADQASAWLT	RQGSFQGGFR
STQDTVIALD	ALSAYWIASH	TTEERGLNVT	LSSTGR	

## [ IDENTIFICATION ]



# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.